

TRACHEOSTOMY RELATED BLEEDING

Staff this document applies to:

Medical staff, Nursing, Physiotherapists and Speech Pathologists on all Austin Health sites

State any related Austin Health policies, procedures or guidelines:

[Emergency Tracheostomy Management Poster](#)

[Suctioning via the tracheostomy](#)

[Tracheostomy Stoma Care](#)

[Medical emergency team \(MET\) call](#)

[Medical emergency review \(MER\) call](#)

[Management of patients with a Tracheostomy](#)

[Mandatory Tracheostomy Equipment](#)

[Escalation Response to Clinical Deterioration](#)

Background:

Tracheostomy is a common procedure involving the placement of an artificial subglottic airway. It may be performed for a variety of reasons, including as part of the management of head & neck cancers, respiratory failure or neurological problems. While serious bleeding from a tracheostomy site is relatively rare, there are occasions where life-threatening erosion into a major vessel can occur. This document is intended to guide the approach to assessing and managing patients with tracheostomy related bleeding so that this problem may be recognised and managed in a timely manner.

Categorising Tracheostomy Related Bleeding:

The potential causes for tracheostomy related bleeding depend very much on the time that has passed since the formation of stoma and tracheostomy tube inserted.

Early bleeding (<4 days)

- Skin related bleeding
- Thyroid related bleeding
- Related to anticoagulant or antiplatelet therapy

Late bleeding (>4) days)

- Erosion into a large artery (e.g. tracheo-innominate fistula)
- Granulation tissue
- Mucosal trauma from suction catheters etc

Response to Tracheostomy Related Bleeding

All bleeding from a tracheostomy site is potentially serious. An experienced clinician should be involved in evaluation at an early stage in all cases. The specialist surgeon or intensivist who inserted the tracheostomy should be consulted immediately and provide advice.

The home team must also be notified of the bleed and also the measures taken.

Recommended action:


- Sit the patient up
- Administer supplemental oxygen
- Measure vital signs
- If actively bleeding or more than 10 millilitres of bright blood is evident (e.g. stoma dressing soaked and blood leaking beyond dressing), activate emergency response e.g. MET/Code Blue.
- If less than 10 millilitres of bright blood (e.g. blood contained within dressing) activate 'Urgent Clinical Review'
- Notify the surgeon responsible for the insertion of the tracheostomy
- In cases of major bleeding and/or the patient has associated hypoxaemia or respiratory distress, call a RESPOND BLUE

Strategies which may temporarily help in the situation of a major arterial erosion include:


- **hyper-inflation** of the tracheostomy **cuff** and/or
- **direct digital compression** of the bleeding point.
- In these circumstances, the patient must have their airway secured by a clinician with advanced airway skills and be transferred to the operating room or interventional radiology suite for further evaluation and intervention. Planning for further investigations and interventions will be at the direction of the treating thoracic, ENT surgeon or Maxillofacial Surgeon involved.
- Ensure adequate wide-bore iv access
- Contact TRAMS by phone extension 3095 or on pager 1291 to report the event. TRAMS will review at the earliest possible opportunity (with the exception of acute ENT or ICU patients which are not managed by TRAMS)

Clinical Alert:

- Small volume bleeding at a tracheostomy stoma may herald a major haemorrhage and the treating clinician must always thoroughly evaluate for the possibility of a trachea-arterial fistula. Please activate the 'urgent clinical review' process
- Tracheo-innominate artery erosion is a rare late complication associated with high mortality rate. Erosion occurs in less than 1% of tracheostomy cases and is usually associated with a) low placement of tracheostomy tube; b) excessive movement of the TT; c) over inflation of cuff; and/or d) suboptimal tracheostomy tube position.
- A pulsating tracheostomy tube may indicate close proximity to a large vessel, which in itself is a risk factor for a major haemorrhage. This constitutes an emergency the first time it is detected and an experienced clinician should be involved in evaluation at an early stage in all cases. The specialist surgeon or intensivist who inserted the tracheostomy should be consulted immediately and provide advice.
- If cuff is inflated, do not deflate cuff until expert clinical assistance is available as it is possible the inflated cuff is tamponading the blood vessel.



Innominate Artery



Implementation Strategy:

All heads of department will be notified via email
 Referred to in all TRAMS education
 Policy will be listed on the TRAMS intranet site and available via ePPIC

Author/Contributors:

Dr Steven Warrillow, Deputy Director, Intensive Care Unit
 Dr James Ward, Respiratory and Sleep Physician, TRAMS Medical Consultant
 Kristy McMurray, TRAMS CNC

Legislation/References/Supporting Documents:

Gray M.C., Mohan S. G, Suxena A., Selvakumar S. The role of innominate artery ligation in the management of massive haemorrhage from tracheo-innominate artery fistula Anaesthesia and Intensive Care Mar 2014;42 266-267
 Grant CA, Dempsey G, Harrison J, Jones T. Tracheo-innominate artery fistula after percutaneous tracheostomy: three case reports and a clinical review. Br J Anaesth 2006;96:127–31
 Seidman, P A; Goldenberg, D and Sinz, E H (eds) Tracheostomy Management: A Multidisciplinary Approach; (2011) Cambridge University Press

Authorised/Endorsed by:

Dr Graeme Hart, Director Intensive Care Unit; TRAMS Team; Dr Mark Howard Director Victorian Respiratory Support Service; Mr Matthew Campbell, Director ENT Service; Mr Simon Knight, Director Thoracic Surgery; Mr Nicholas Rutherford, Director Maxillofacial Surgery; Larry McNichol CSU Medical Director – Anaesthesia, Perioperative and Intensive Care CSU; Nursing Standards Committee; Clinical Policy and Procedures Committee

Primary Person/Department Responsible for Document:

Dr James Ward, Respiratory and Sleep Physician, TRAMS Medical Consultant